


Recommended immunizations for children from birth through 6 years old*



NOTE: If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have questions about vaccines.

FOOTNOTES: * The recommended immunization chart shown above was updated in 2014.

[†] Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

[‡] Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

If your child has any medical conditions that put him at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.

For more information, call toll free
1-800-CDC-INFO (1-800-232-4636)
or visit
<http://www.cdc.gov/vaccines>



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

**AMERICAN ACADEMY OF
FAMILY PHYSICIANS**
STRONG MEDICINE FOR AMERICA

**American Academy
of Pediatrics**



DEDICATED TO THE HEALTH OF ALL CHILDREN™

DTaP=diphtheria, tetanus, and pertussis; HepA=hepatitis A; HepB=hepatitis B; Hib=Haemophilus influenzae type b; IPV=inactivated polio vaccine; MMR=measles, mumps, and rubella; PCV=pneumococcal conjugate vaccine; RV=rotavirus vaccine.
This content was adapted by Pfizer from information provided by the Centers for Disease Control and Prevention.